


# SteatoTest

Internal Reference : FAKE TEST JMC

Demographics	
Sex	Male
Birthdate	16/08/1977

Analysis		
Sample Date	07/04/2006	
Alpha2 Macroglobulin (g/l)	1.20	
Haptoglobin (g/l)	1.10	
Apolipoprotein A1 (g/l)	0.90	
Bilirubin (µMol/l)	22.00	
Gamma GT (IU/l)	66	
ALT (IU/l)	50	
AST (IU/l)	66	
Fasting glucose (mmol/l)	4.00	
Triglycerides (mmol/l)	1.10	
Total cholesterol (mmol/l)	4.00	
Weight (kg)	77.00	
Height (meters)	1.70	

## SteatoTest



Score: 0.59  
(S2)

### Usage precautions

- The reliability of results is bound with the respect of the preanalytical and analytical conditions recommended by BioPredictive.
- The Tests have to be deferred for: acute hemolysis, acute hepatitis, acute inflammation, extra hepatic cholestasis.
- The advice of a specialist should be sought for interpretation in chronic hemolysis and Gilbert's syndrome.
- The Test interpretation is not validated in liver transplant patients.
- Isolated extreme values of one of the components should lead to caution in interpreting the results.
- In case of discordance between a biopsy result and a Test, it is advised to seek for the advice of a specialist. The causes of these discordances could be due to a flaw of the Test or to a flaw in the biopsy: i.e. a liver biopsy has a 33% variability rate for one fibrosis stage
- FibroTest and SteatoTest are interpretable for chronic hepatitis B and C, alcoholic and non alcoholic steatosis.
- NashTest is interpretable for non alcoholic steatosis.
- AshTest is interpretable for alcoholic steatosis.
- ActiTest is interpretable for chronic hepatitis B and C.